

DISTRIBUTIONS OF DISCRIMINANTS OF CUBIC ALGEBRAS II

TAKASHI TANIGUCHI

ABSTRACT. Let k be a number field and \mathcal{O} the ring of integers. In the previous paper [T06] we study the Dirichlet series counting discriminants of cubic algebras of \mathcal{O} and derive some density theorems on distributions of the discriminants by using the theory of zeta functions of prehomogeneous vector spaces. In this paper we consider these objects under imposing finite number of splitting conditions at non-archimedean places. Especially the explicit formulae of residues at $s = 1$ and $5/6$ under the conditions are given.

1. INTRODUCTION

Let k be a number field and \mathcal{O} the ring of integers. For a place v of k let k_v be the completion of k at v . Let T be a finite set of places. Take a separable cubic algebra L_v of k_v for each $v \in T$ and let $L_T = (L_v)_{v \in T}$ the T -tuple. We call an \mathcal{O} -algebra a *cubic algebra* if it is locally free of rank 3 as an \mathcal{O} -module. We denote by $\mathcal{C}(\mathcal{O})$ the set of isomorphism classes of cubic algebras of \mathcal{O} . We let

$$\mathcal{C}(\mathcal{O}, L_T)^{\text{ird}} := \left\{ R \in \mathcal{C}(\mathcal{O}) \mid \begin{array}{l} F = R \otimes_{\mathcal{O}} k \text{ is a cubic field extension of } k, \text{ and} \\ F \otimes_k k_v \cong L_v \text{ for all } v \in T. \end{array} \right\}.$$

We define

$$\vartheta_{L_T}^{\text{ird}}(s) := \sum_{R \in \mathcal{C}(\mathcal{O}, L_T)^{\text{ird}}} \frac{\#(\text{Aut}(R))^{-1}}{N(\Delta_{R/\mathcal{O}})^s},$$

$$h_{L_T}(X) := \# \{ R \in \mathcal{C}(\mathcal{O}, L_T)^{\text{ird}} \mid N(\Delta_{R/\mathcal{O}}) < X \}.$$

Here we denote by $N(\Delta_{R/\mathcal{O}})$ the ideal norm of the relative discriminant of R/\mathcal{O} and by $\text{Aut}(R)$ the group of automorphisms of R as an \mathcal{O} -algebra. The primary purpose of this paper is to prove the following. Let $n = [k : \mathbb{Q}]$.

Theorem 1.1. *There exist constants \mathfrak{A}_{L_T} and \mathfrak{B}_{L_T} described explicitly such that;*

- (1) $\vartheta_{L_T}^{\text{ird}}(s)$ has meromorphic continuation to the whole complex plane which is holomorphic for $\text{Re}(s) > 1/2$ except for simple poles at $s = 1$ and $5/6$ with residues \mathfrak{A}_{L_T} and \mathfrak{B}_{L_T} , respectively, and
- (2) for any $\varepsilon > 0$,

$$h_{L_T}(X) = \mathfrak{A}_{L_T} X + (5/6)^{-1} \mathfrak{B}_{L_T} X^{5/6} + O(X^{\frac{5n-1}{5n+1} + \varepsilon}) \quad (X \rightarrow \infty).$$

Note that the $X^{5/6}$ -term in the formula is relevant only when $n = 1, 2$. We give the formulae of \mathfrak{A}_{L_T} and \mathfrak{B}_{L_T} . Let $\zeta_k(s)$ and Δ_k be the Dedekind zeta function and the absolute discriminant of k , respectively. We denote by \mathfrak{M}_{∞} , $\mathfrak{M}_{\mathbb{R}}$, $\mathfrak{M}_{\mathbb{C}}$ and \mathfrak{M}_{f} the set of all infinite places, real places, complex places and finite places, respectively. We put $r_1 = \#\mathfrak{M}_{\mathbb{R}}$ and $r_2 = \#\mathfrak{M}_{\mathbb{C}}$. For $v \in \mathfrak{M}_{\text{f}}$, let q_v be the order of the residue field of k_v . We put $\theta_{L_v} = \#(\text{Aut}_{k_v\text{-algebra}}(L_v))$. For a non-archimedean local field K with the order of residue field q , we define its local zeta function by $\zeta_K(s) = (1 - q^{-s})^{-1}$. The cubic algebra L_v is in general a product of local fields. We define

Date: February 2, 2008.

Financial support is provided by Research Fellowships for Young Scientists of Japan Society for the Promotion of Science.

$\zeta_{L_v}(s)$ as the product of the zeta functions of those fields. The relative discriminant Δ_{L_v/k_v} is also defined as the product of relative discriminants of those local fields. We denote by Δ_{L_v} its norm. We put $i_\infty(L_T) = \#\{v \in \mathfrak{M}_\mathbb{R} \mid L_v = \mathbb{R}^3\}$. We give the value in case of $T \supset \mathfrak{M}_\infty$. The general case is easily obtained from this by taking a suitable summation.

Theorem 1.2. *When $T \supset \mathfrak{M}_\infty$, the constants \mathfrak{A}_{L_T} and \mathfrak{B}_{L_T} are given by*

$$\mathfrak{A}_{L_T} = \frac{\text{Res}_{s=1} \zeta_k(s) \cdot \zeta_k(2)}{2^{r_1+r_2+1} 3^{i_\infty(L_T)+r_2}} \prod_{v \in T \cap \mathfrak{M}_f} \alpha_v(L_v),$$

$$\mathfrak{B}_{L_T} = \frac{\text{Res}_{s=1} \zeta_k(s) \cdot \zeta_k(1/3)}{6 \Delta_k^{1/2} (\sqrt{3})^{r_2-i_\infty(L_T)}} \left(\frac{3\Gamma(1/3)^3}{2\pi} \right)^n \prod_{v \in T \cap \mathfrak{M}_f} \beta_v(L_v),$$

where

$$\alpha_v(L_v) = \frac{(1 - q_v^{-1})(1 - q_v^{-2})}{(1 - q_v^{-4})(1 - q_v^{-5})} \cdot \theta_{L_v}^{-1} \Delta_{L_v}^{-1} \cdot \frac{\zeta_{L_v}(2)}{\zeta_{L_v}(4)},$$

$$\beta_v(L_v) = \frac{(1 - q_v^{-1/3})(1 - q_v^{-1})}{(1 - q_v^{-10/3})(1 - q_v^{-4})} \cdot \theta_{L_v}^{-1} \Delta_{L_v}^{-1} \cdot \frac{\zeta_{L_v}(1/3) \zeta_{L_v}(5/3)}{\zeta_{L_v}(2/3) \zeta_{L_v}(10/3)}.$$

Let $v \in \mathfrak{M}_f$. We see by computation that $\sum_{L_v} \alpha_v(L_v) = \sum_{L_v} \beta_v(L_v) = 1$ where L_v runs through all the separable cubic algebras of k_v . Hence $\alpha_v(L_v)$ and $\beta_v(L_v)$ give the proportion of the contributions of cubic algebras with local splitting type L_v . The computation of $\alpha_v(L_v)$ is reduced to the determination of certain orbital volume in a v -adic vector space. The meaning of $\beta_v(L_v)$ is more subtle and the computation requires a careful local theory.

In this paper we also try to improve Theorem 1.1 under segregating the cubic algebras via the *Steinitz class*. The Steinitz class takes value in the ideal class group, and it is known that finitely generated locally free modules over a Dedekind domain is completely classified by the rank and the Steinitz class (cf. [M71].) Let $\text{Cl}(k)$ be the ideal class group of k . For $R \in \mathcal{C}(\mathcal{O})$, let $\text{St}(R) \in \text{Cl}(k)$ be its Steinitz class. This is the class of a fractional ideal isomorphic to $\bigwedge^3 R$ as an \mathcal{O} -module. For $\mathfrak{a} \in \text{Cl}(k)$, we define

$$\vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s) := \sum_{R \in \mathcal{C}(\mathcal{O}, L_T)^{\text{ird}}, \text{St}(R) = \mathfrak{a}} \frac{\#(\text{Aut}(R))^{-1}}{N(\Delta_{R/\mathcal{O}})^s},$$

$$h_{L_T}(\mathfrak{a}, X) := \#\{R \in \mathcal{C}(\mathcal{O}, L_T)^{\text{ird}} \mid \text{St}(R) = \mathfrak{a}, N(\Delta_{R/\mathcal{O}}) < X\}.$$

Then obviously $\vartheta_{L_T}^{\text{ird}}(s) = \sum_{\mathfrak{a} \in \text{Cl}(k)} \vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s)$ and $h_{L_T}(X) = \sum_{\mathfrak{a} \in \text{Cl}(k)} h_{L_T}(\mathfrak{a}, X)$. Let h_k be the class number of k and $h_k^{(3)}$ the number of 3-torsions of $\text{Cl}(k)$ (which is a power of 3.) Also for $\mathfrak{a} \in \text{Cl}(k)$, we put $\tau(\mathfrak{a}) = 1$ if there exists $\mathfrak{b} \in \text{Cl}(k)$ such that $\mathfrak{a} = \mathfrak{b}^3$ and $\tau(\mathfrak{a}) = 0$ otherwise. In this paper we also prove the following.

Theorem 1.3. (1) *The Dirichlet series $\vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s)$ has a meromorphic continuation to the whole complex plane which is holomorphic for $\text{Re}(s) > 1/2$ except for a simple poles at $s = 1$ with the residue \mathfrak{A}_{L_T}/h_k and a possible simple pole at $s = 5/6$ with the residue $\tau(\mathfrak{a}) \mathfrak{B}_{L_T} h_k^{(3)}/h_k$, respectively.*

(2) *Assume L_T is chosen so that at least one of L_v is a field. Then $\vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s)$ is holomorphic in the whole complex plane except for $s = 1$ and $5/6$. Also for any $\varepsilon > 0$,*

$$h_{L_T}(\mathfrak{a}, X) = \frac{\mathfrak{A}_{L_T}}{h_k} X + \tau(\mathfrak{a}) \frac{h_k^{(3)}}{h_k} \cdot \frac{\mathfrak{B}_{L_T}}{5/6} X^{5/6} + O(X^{\frac{4n-1}{4n+1} + \varepsilon}) \quad (X \rightarrow \infty).$$

The statement (1) is an improvement of Theorem 1.1 (1). In case $[k : \mathbb{Q}] = 2$ and $3 \mid h_k$, the formula in (2) implies that the distribution of the Steinitz classes of the elements of $\mathcal{C}(\mathcal{O}, L_T)^{\text{ird}}$

is not uniform, and the irregularity is reflected in the $X^{5/6}$ -term. Although it is likely that such a formula exists for unconditional L_T also, to remove the condition on L_T is highly non-trivial.

In the previous paper [T06] we prove these theorems for $T = \mathfrak{M}_\infty$. (The original Shintani's theorem [S75, Theorem 4] is for $k = \mathbb{Q}$, $T = \{\infty\}$.) As in [T06] we approach these theorems by using *Sato-Shintani's zeta function* [SS74] for the space of binary cubic forms.

The study of class numbers of integral binary cubic forms over \mathbb{Z} was initiated by G. Eisenstein and developed by many mathematicians including C. Hermite, H. Davenport and T. Shintani. Via Delone-Faddeev's correspondence [DF64] the $\mathrm{GL}_2(\mathbb{Z})$ -orbits of integral binary cubic forms corresponds bijectively to the set of cubic rings, thus useful to investigate cubic fields or their orders. Shintani [S72] introduced Dirichlet series whose coefficients are the class numbers and studied extensively as an example of *zeta functions of prehomogeneous vector spaces*. This was generalized to over a general number field using adelic language in [W85], and used to investigate relative cubic extensions over the base field ([DW86, DW88]) or its integer ring ([T06].) Besides the rightmost pole at $s = 1$, the Dirichlet series has a mysterious second pole at $s = 5/6$. Our purpose is to study how those residues behave when local conditions at finite places are imposed.

We prove theorems above in the following process. In Section 2 we review the notation and invariant measure of [T06] those we use in this paper. In Section 3 we recall and refine the parameterizations of cubic algebras by means of the space of binary cubic form that we established in the previous paper [T06]. In Section 4 we introduce the global zeta function for the space of binary cubic forms. We also introduce two partial zeta integrals, those the contributions from irreducible forms and from reducible forms. We use the result of Section 3 to express those integrals by Dirichlet series counting cubic algebras.

In Section 5 we give the analytic continuations and residue formulae of some Dirichlet series including $\vartheta_{L_T}^{\mathrm{ird}}(\mathfrak{a}, s)$. This give a proof of Theorem 1.1 (1) and Theorem 1.3 (1) with the values \mathfrak{A}_{L_T} , \mathfrak{B}_{L_T} in Theorem 1.2. The meromorphic continuations and residues of the zeta integrals are obtained in [T06, Section 8]. The residues are expressed by means of local distributions and hence the computation of the residues are reduced to the local theory. The local theory for the space of binary cubic forms were studied by Datskovsky-Wright [DW86] in detail, and their results make us the computation simple. We note that the explicit formula [DW86, Theorem 3.1] of the non-archimedean local zeta function plays an important role.

In Section 6 we prove Theorem 1.1 (2) and Theorem 1.3 (2). Our tool to find density theorems is a modified version [SS74, Theorem 3] of Landau's Tauberian theorem [L15, Hauptsatz], using a functional equation to derive some informations on the error term. To separate the contributions of reducible forms, we also use the functional equations of zeta functions for the space of binary quadratic forms.

2. NOTATION AND INVARIANT MEASURES

For notation we basically follow [T06]. But we have one exception. If V is a scheme define over a ring R and S is an R -algebra, then in this paper we denote the set of S -rational points of V by $V(S)$, not by V_S as we denoted in [T06].

For a finite set X we denote by $\#X$ its cardinality. If an abstract group G acts on a set X , then for $x \in X$ we set $\mathrm{Stab}(G; x) = \{g \in G \mid gx = x\}$. If $\mathfrak{x} \in G \backslash X$ is the class of $x \in X$, we also denote the group by $\mathrm{Stab}(G; \mathfrak{x})$, which is well defined up to isomorphism. The one-dimensional affine space is denoted by Aff .

Throughout this paper we fix a number field k . We use the notation \mathcal{O} , n , $\zeta_k(s)$, Δ_k , \mathfrak{M}_∞ , $\mathfrak{M}_{\mathbb{R}}$, $\mathfrak{M}_{\mathbb{C}}$, $\mathfrak{M}_{\mathfrak{f}}$, r_1 , r_2 , k_v , q_v , $\mathrm{Cl}(k)$, h_k , $h_k^{(3)}$ and $\tau(\mathfrak{a})$ as in Section 1. Let \mathfrak{M} be the set of all places of k . We put $\mathfrak{c}_k = \mathrm{Res}_{s=1} \zeta_k(s)$. For a fractional ideal I of k , we denote by $N(I)$ its ideal norm. The rings of adeles and finite adeles are denoted by \mathbb{A} and $\mathbb{A}_{\mathfrak{f}}$. We put $k_\infty = k \otimes_{\mathbb{Q}} \mathbb{R}$ and $\widehat{\mathcal{O}} = \mathcal{O} \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}}$. Note that $\widehat{\mathcal{O}} = \prod_{v \in \mathfrak{M}_{\mathfrak{f}}} \mathcal{O}_v$, $\mathbb{A}_{\mathfrak{f}} = \widehat{\mathcal{O}} \otimes_{\mathcal{O}} k$, and $\mathbb{A} = k_\infty \times \mathbb{A}_{\mathfrak{f}}$. The adelic

absolute value $|\cdot|_{\mathbb{A}}$ on \mathbb{A}^\times is normalized so that, for $t \in \mathbb{A}^\times$, $|t|_{\mathbb{A}}$ is the module of multiplication by t with respect to any Haar measure dx on \mathbb{A} , i.e. $|t|_{\mathbb{A}} = d(tx)/dx$. We define $|\cdot|_\infty$, $|\cdot|_{\mathbb{A}_f}$ and $|\cdot|_v$ on k_∞^\times , \mathbb{A}_f^\times and k_v^\times similarly. For a vector space V , Let $\mathcal{S}(V(\mathbb{A}))$, $\mathcal{S}(V(k_v))$, $\mathcal{S}(V(k_\infty))$ and $\mathcal{S}(V(\mathbb{A}_f))$ be the spaces of Schwartz–Bruhat functions on each of the indicated domains.

For a fractional ideal \mathfrak{a} , let $i(\mathfrak{a}) \in \mathbb{A}_f^\times (\subset \mathbb{A}^\times)$ be the corresponding idele, which is well defined up to $\widehat{\mathcal{O}}^\times$ -multiple. That is, $i(\mathfrak{a}) \in \mathbb{A}_f^\times$ is characterized by the condition $\mathfrak{a} = k \cap i(\mathfrak{a})\widehat{\mathcal{O}}$. Then $|i(\mathfrak{a})|_{\mathbb{A}} = N(\mathfrak{a})^{-1}$. Notice that the infinite component of $i(\mathfrak{a})$ is trivial. If there is no confusion we simply write \mathfrak{a} instead of $i(\mathfrak{a})$. The set of characters of $\text{Cl}(k)$ is denoted by $\text{Cl}(k)^*$. We regard $\omega \in \text{Cl}(k)^*$ as a character on $\mathbb{A}^\times/k^\times$ via the standard composition of the maps $\mathbb{A}^\times/k^\times \rightarrow \mathbb{A}^\times/k_\infty^\times k^\times \widehat{\mathcal{O}}^\times \cong \text{Cl}(k) \rightarrow \mathbb{C}^\times$. Then $\omega(\mathfrak{a}) = \omega(i(\mathfrak{a}))$.

We give normalizations of invariant measures. In general, for an algebraic group X over k with local measures dx_v on $X(k_v)$ for all $v \in \mathfrak{M}$ are given, then we always denote by $dx_\infty = \prod_{v \in \mathfrak{M}_\infty} dx_v$, $dx_f = \prod_{v \in \mathfrak{M}_f} dx_v$ and $d_{\text{pr}x} = \prod_{v \in \mathfrak{M}} dx_v$, which are measures on $X(k_\infty)$, $X(\mathbb{A}_f)$ and $X(\mathbb{A})$, respectively.

For any $v \in \mathfrak{M}_f$, we choose a Haar measure dx_v on k_v to satisfy $\int_{\mathcal{O}_v} dx_v = 1$. We write dx_v for the ordinary Lebesgue measure if v is real, and for twice the Lebesgue measure if v is imaginary. For any $v \in \mathfrak{M}_f$, we normalize the Haar measure $d^\times t_v$ on k_v^\times such that $\int_{\mathcal{O}_v^\times} d^\times t_v = 1$. Let $d^\times t_v(x) = |x|_v^{-1} dx_v$ if $v \in \mathfrak{M}_\infty$.

Let $G = \text{GL}_2$. We review the normalization of the measure on $G(k_v) = \text{GL}_2(k_v)$. Let $B \subset G$ be the Borel subgroup consisting of lower triangular matrices. We normalize the right invariant measure db_v on $B(k_v)$ by

$$\int_{B_{k_v}} f(b_v) db_v = \int_{(k_v^\times)^2 \times k_v} f\left(\begin{pmatrix} s_v & 0 \\ 0 & t_v \end{pmatrix} \begin{pmatrix} 1 & 0 \\ u_v & 1 \end{pmatrix}\right) \left|\frac{t_v}{s_v}\right|_v d^\times s_v d^\times t_v du_v.$$

Let $\mathcal{K} = \prod_{v \in \mathfrak{M}} \mathcal{K}_v$ where $\mathcal{K}_v = \text{O}(2), \text{U}(2), \text{GL}_2(\mathcal{O}_v)$ for $v \in \mathfrak{M}_\mathbb{R}, \mathfrak{M}_\mathbb{C}, \mathfrak{M}_f$, respectively. We choose an invariant measure $d\kappa_v$ on \mathcal{K}_v such that $\int_{\mathcal{K}_v} d\kappa_v = 1$. The group $G(k_v)$ has the decomposition $G(k_v) = \mathcal{K}_v B(k_v)$. We choose an invariant measure on $G(k_v)$ by $dg_v = d\kappa_v db_v$ for $g_v = \kappa_v b_v$.

Finally we express diagonal elements of G as $\text{diag}(t_1, t_2) = \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix}$.

3. THE SPACE OF BINARY CUBIC FORMS AND PARAMETERIZATION

Let G be the general linear group of rank 2 and V the space of binary cubic forms;

$$G = \text{GL}_2,$$

$$V = \{x = x(v_1, v_2) = x_0 v_1^3 + x_1 v_1^2 v_2 + x_2 v_1 v_2^2 + x_3 v_2^3 \mid x_i \in \text{Aff}\}.$$

We define the action of G on V by

$$(gx)(v) = (\det g)^{-1} x(vg).$$

The twist by $\det(g)^{-1}$ is to make the representation faithful. For $x \in V$, let $P(x)$ be the discriminant;

$$P(x) = x_1^2 x_2^2 - 4x_0 x_2^3 - 4x_1^3 x_3 + 18x_0 x_1 x_2 x_3 - 27x_0^2 x_3^2.$$

Then we have $P(gx) = (\det g)^2 P(x)$. We put $V^{\text{ss}} = \{x \in V \mid P(x) \neq 0\}$.

We first recall the parameterization of cubic algebras of \mathcal{O} . Let \mathfrak{a} be a fractional ideal of k . We put

$$G(k) \supset G_{\mathfrak{a}} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a \in \mathcal{O}, b \in \mathfrak{a}, c \in \mathfrak{a}^{-1}, d \in \mathcal{O}, ad - bc \in \mathcal{O}^\times \right\},$$

$$V(k) \supset V_{\mathfrak{a}} = \{x \mid x_0 \in \mathfrak{a}, x_1 \in \mathcal{O}, x_2 \in \mathfrak{a}^{-1}, x_3 \in \mathfrak{a}^{-2}\}.$$

Then $V_{\mathfrak{a}}$ is a $G_{\mathfrak{a}}$ -invariant submodule. As in Section 1, let $\mathcal{C}(\mathcal{O})$ be the set of isomorphism classes of cubic algebras of \mathcal{O} . For $R \in \mathcal{C}(\mathcal{O})$, let $\text{Aut}(R)$ be the group of automorphisms of R as an \mathcal{O} -algebra. We put $\mathcal{C}(\mathcal{O}, \mathfrak{a}) = \{R \in \mathcal{C}(\mathcal{O}) \mid \text{St}(R) = \mathfrak{a}\}$. In [T06, Section 3] we establish the following parameterization of $\mathcal{C}(\mathcal{O}, \mathfrak{a})$.

Proposition 3.1. (1) *There exists the canonical bijection between $\mathcal{C}(\mathcal{O}, \mathfrak{a})$ and $G_{\mathfrak{a}} \backslash V_{\mathfrak{a}}$ making the following diagram commutative:*

$$\begin{array}{ccc} G_{\mathfrak{a}} \backslash V_{\mathfrak{a}} & \xrightarrow{\quad} & \mathcal{C}(\mathcal{O}, \mathfrak{a}) \\ \downarrow P & & \downarrow \text{discriminant} \\ (\mathcal{O}^{\times})^2 \backslash \mathfrak{a}^{-2} & \xrightarrow{\times \mathfrak{a}^2} & \{\text{integral ideals of } \mathcal{O}\}. \end{array}$$

Here, the right vertical arrow is to take the discriminant, and the low horizontal arrow is given by multiplying \mathfrak{a}^2 . For each $R \in \mathcal{C}(\mathcal{O}, \mathfrak{a})$, we denote by x_R the corresponding element in $G_{\mathfrak{a}} \backslash V_{\mathfrak{a}}$ or its arbitrary representative in $V_{\mathfrak{a}}$.

(2) *We have $\text{Aut}(R) \cong \text{Stab}(G_{\mathfrak{a}}; x_R)$.*

Let K be either the number field k or a local field k_v . We next recall the geometric interpretation of the $G(K)$ -orbits in $V^{\text{ss}}(K)$. We denote by $\mathcal{C}^{\text{sep}}(K)$ the set of isomorphism classes of separable cubic algebras of K . For $x = x(v_1, v_2) \in V^{\text{ss}}(K)$, we define

$$Z_x = \text{Proj } K[v_1, v_2]/(x(v_1, v_2)), \quad K(x) = \Gamma(Z_x, \mathcal{O}_{Z_x}).$$

We regard $K(x)$ as an element of $\mathcal{C}^{\text{sep}}(K)$. It is well known that The map $x \mapsto K(x)$ gives a bijection between $G(K) \backslash V^{\text{ss}}(K)$ and $\mathcal{C}^{\text{sep}}(K)$. By definition, this correspondence is functorial with respect to the localization, i.e., for $x \in V^{\text{ss}}(k)$ and $v \in \mathfrak{M}$ we have $k_v(x) = k(x) \otimes_k k_v$. We have the following compatibility of the two parameterizations $x \mapsto x_R$ and $x \mapsto k(x)$.

Lemma 3.2. *Let $R \in \mathcal{C}(\mathcal{O}, \mathfrak{a})$ and $\Delta_{R/\mathcal{O}} \neq 0$. Then $x_R \in V^{\text{ss}}(k)$ and $k(x_R) = R \otimes_{\mathcal{O}} k$.*

For $L_v \in \mathcal{C}^{\text{sep}}(k_v)$, we put

$$V_{L_v} = \{x \in V^{\text{ss}}(k_v) \mid k_v(x) = L_v\},$$

which is the $G(k_v)$ -orbit in $V^{\text{ss}}(k_v)$ corresponding to L_v . This *should not be confused* to the set of L_v -rational points of V nor the base change of V to L_v . Since we don't consider such objects in this paper we hope this notation is not misleading.

Definition 3.3. Let T be a set of places. Let $L_T = (L_v)_{v \in T}$ be a T -tuple where $L_v \in \mathcal{C}^{\text{sep}}(k_v)$ for each $v \in T$. For such L_T , we define

$$\begin{aligned} V_{k, L_T} &= \{x \in V^{\text{ss}}(k) \mid k(x) \otimes_k k_v = L_v \text{ for all } v \in T\}, \\ \mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T) &= \{R \in \mathcal{C}(\mathcal{O}) \mid \text{St}(R) = \mathfrak{a}, R \otimes_{\mathcal{O}} k_v = L_v \text{ for all } v \in T\}. \end{aligned}$$

Proposition 3.4. *The correspondence in Proposition 3.1 induces a bijection between $G_{\mathfrak{a}} \backslash (V_{\mathfrak{a}} \cap V_{k, L_T})$ and $\mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T)$.*

Proof. Let $R \in \mathcal{C}(\mathcal{O}, \mathfrak{a})$. By Lemma 3.2, $R \otimes_{\mathcal{O}} k_v = k(x_R) \otimes_k k_v$. Hence $R \otimes_{\mathcal{O}} k_v = L_v$ if and only if $x_R \in V_{L_v}$. Since $V_{k, L_T} = \bigcap_{v \in T} (V^{\text{ss}}(k) \cap V_{L_v})$, we have the bijection. \square

Let $V^{\text{ss}}(k)^{\text{ird}}$ (resp. $V^{\text{ss}}(k)^{\text{rd}}$) be the subset of $V^{\text{ss}}(k)$ consisting of binary cubic forms irreducible (resp. reducible) over k . By the definition of $k(x)$, $V^{\text{ss}}(k)^{\text{ird}} = \{x \in V^{\text{ss}}(k) \mid k(x)/k \text{ is a cubic field extension}\}$. Also we put

$$\begin{aligned} \mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T)^{\text{ird}} &= \{R \in \mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T) \mid R \otimes_{\mathcal{O}} k \text{ is a cubic field extension of } k\}, \\ \mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T)^{\text{rd}} &= \{R \in \mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T) \mid R \otimes_{\mathcal{O}} k \text{ is a separable algebra and not a field}\}. \end{aligned}$$

Then by Lemma 3.2, we have the following refined version of Proposition 3.4.

Proposition 3.5. *The correspondence of Proposition 3.1 induces a bijection between the following sets;*

- (1) $G_{\mathfrak{a}} \backslash (V_{\mathfrak{a}} \cap V_{k, L_T} \cap V^{\text{ss}}(k)^{\text{ird}})$ and $\mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T)^{\text{ird}}$.
- (2) $G_{\mathfrak{a}} \backslash (V_{\mathfrak{a}} \cap V_{k, L_T} \cap V^{\text{ss}}(k)^{\text{rd}})$ and $\mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T)^{\text{rd}}$.

4. GLOBAL ZETA FUNCTIONS AND DIRICHLET SERIES

In this section we introduce the global zeta function and related zeta integrals. Using Propositions 3.4 and 3.5, we see that these are integral expressions of the Dirichlet series we want to investigate.

Definition 4.1. For $\Phi \in \mathcal{S}(V_{\mathbb{A}})$, $s \in \mathbb{C}$ and $\omega \in \text{Cl}(k)^*$, we define the *global zeta function* by

$$Z(\Phi, s, \omega) = \int_{G(\mathbb{A})/G(k)} |\det g|^{2s} \omega(\det g) \sum_{x \in V^{\text{ss}}(k)} \Phi(gx) d_{\text{pr}} g.$$

Also for $\mathfrak{a} \in \text{Cl}(k)$, we define

$$Z_{\mathfrak{a}}(\Phi, s) = \int_{G(k_{\infty})G(\widehat{\mathcal{O}})\text{diag}(1, \mathfrak{a})G(k)/G(k)} |\det g|^{2s} \sum_{x \in V^{\text{ss}}(k)} \Phi(gx) d_{\text{pr}} g.$$

Recall the double coset decomposition $G(\mathbb{A}) = \coprod_{\mathfrak{a} \in \text{Cl}(k)} G(k_{\infty})G(\widehat{\mathcal{O}})\text{diag}(1, \mathfrak{a})G(k)$. Since $\omega(\det(G(k_{\infty})G(\widehat{\mathcal{O}})G(k))) = \omega(k_{\infty}^{\times} \widehat{\mathcal{O}}^{\times} k^{\times}) = 1$, we have the following.

Lemma 4.2.

$$Z(\Phi, s, \omega) = \sum_{\mathfrak{a} \in \text{Cl}(k)} \omega(\mathfrak{a}) Z_{\mathfrak{a}}(\Phi, s).$$

Let $T \supset \mathfrak{M}_{\infty}$ be a finite set of places. For the rest of this paper we fix T and $L_v \in \mathcal{C}^{\text{sep}}(k_v)$ for each $v \in T$. We put $L_T = (L_v)_{v \in T}$. For our purpose, we also assume $\Phi \in \mathcal{S}(V(\mathbb{A}))$ is chosen as follows for the rest of this section.

Assumption 4.3. We assume Φ is the product $\Phi = \prod_{v \in \mathfrak{M}} \Phi_v$, where

$$\Phi_v \text{ is } \begin{cases} \text{an arbitrary } \mathcal{K}_v\text{-invariant function supported in } V_{L_v} & v \in \mathfrak{M}_{\infty}, \\ \text{the characteristic function of } V(\mathcal{O}_v) \cap V_{L_v} & v \in \mathfrak{M}_f \cap T, \\ \text{the characteristic function of } V(\mathcal{O}_v) & v \in \mathfrak{M}_f \setminus T. \end{cases}$$

We put $\Phi_{\infty} = \prod_{v \in \mathfrak{M}_{\infty}} \Phi_v$ and $\Phi_f = \prod_{v \in \mathfrak{M}_f} \Phi_v$. Note that each Φ_v is $G(\mathcal{O}_v)$ -invariant for $v \in \mathfrak{M}_f$ and hence Φ_f is $G(\widehat{\mathcal{O}})$ -invariant.

Definition 4.4. We define

$$\begin{aligned} \mathcal{Z}_{L_T}(\Phi_{\infty}, s) &= \int_{G(k_{\infty})} |P(g_{\infty}x)|_{\infty}^s \Phi_{\infty}(g_{\infty}x) dg_{\infty} \quad (x \in V_{k, L_T}), \\ \vartheta_{L_T}(\mathfrak{a}, s) &= \sum_{x \in G_{\mathfrak{a}} \backslash (V_{\mathfrak{a}} \cap V_{k, L_T})} \frac{(\#(\text{Stab}(G_{\mathfrak{a}}; x)))^{-1}}{N(\mathfrak{a})^{2s} |P(x)|_{\infty}^s} = \sum_{R \in \mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T)} \frac{(\# \text{Aut}(R))^{-1}}{N(\Delta_R/\mathcal{O})^s}. \end{aligned}$$

The function $\mathcal{Z}_{L_T}(\Phi_{\infty}, s)$ is called the *local zeta function*. Since $V_{k, L_T} \subset V(k_{\infty})$ is contained in a single $G(k_{\infty})$ -orbit $\prod_{v \in \mathfrak{M}_{\infty}} V_{L_v}$, this does not depend on the choice of x . Note that the second equality in the lower formula follows from Propositions 3.1 and 3.4.

Lemma 4.5.

$$Z_{\mathfrak{a}}(\Phi, s) = \mathcal{Z}_{L_T}(\Phi_{\infty}, s) \vartheta_{L_T}(\mathfrak{a}, s).$$

Proof. Let $G(\widehat{\mathcal{O}})_{\mathfrak{a}} = \text{diag}(1, \mathfrak{a})^{-1} G(\widehat{\mathcal{O}}) \text{diag}(1, \mathfrak{a})$ and $\Phi_{\mathfrak{a}}(x) = \Phi(\text{diag}(1, \mathfrak{a})x)$. Since the infinite part of $\mathfrak{a} \in \mathbb{A}^\times$ is trivial, the infinite part of $\Phi_{\mathfrak{a}}$ coincides with Φ_∞ . We denote by $\Phi_{\mathfrak{f}, \mathfrak{a}}$ the finite part of $\Phi_{\mathfrak{a}}$, hence $\Phi_{\mathfrak{a}} = \Phi_\infty \times \Phi_{\mathfrak{f}, \mathfrak{a}}$. Then $\Phi_{\mathfrak{f}, \mathfrak{a}}$ is $G(\widehat{\mathcal{O}})_{\mathfrak{a}}$ -invariant. Hence we have

$$\begin{aligned} Z_{\mathfrak{a}}(\Phi, s) &= \frac{1}{N(\mathfrak{a})^{2s}} \int_{G(k_\infty)G(\widehat{\mathcal{O}})_{\mathfrak{a}}G(k)/G(k)} |\det g|_{\mathbb{A}}^{2s} \sum_{x \in V^{ss}(k)} \Phi_{\mathfrak{a}}(gx) dg \\ &= \frac{1}{N(\mathfrak{a})^{2s}} \int_{G(k_\infty)G(\widehat{\mathcal{O}})_{\mathfrak{a}}/G(k) \cap G(k_\infty)G(\widehat{\mathcal{O}})_{\mathfrak{a}}} |\det g_\infty|_\infty^{2s} \sum_{x \in V^{ss}(k)} \Phi_{\mathfrak{f}, \mathfrak{a}}(x) \Phi_\infty(g_\infty x) dg_\infty dg_{\mathfrak{f}}. \end{aligned}$$

For $x \in V^{ss}(k)$, $\Phi_{\mathfrak{f}, \mathfrak{a}}(x) = 1$ if

$$\text{diag}(1, \mathfrak{a})x \in V(\widehat{\mathcal{O}}) \quad \text{and} \quad \text{diag}(1, \mathfrak{a})_v x \in V_{L_v} \quad \text{for all } v \in T,$$

and 0 otherwise. Here $\text{diag}(1, \mathfrak{a})_v$ denote the v -component of $\text{diag}(1, \mathfrak{a})$. Since V_{L_v} is a $G(k_v)$ -orbit, the second condition holds precisely when $x \in V_{L_v}$ for all $v \in T$. Also we see $\text{diag}(1, \mathfrak{a})^{-1}V(\widehat{\mathcal{O}}) \cap V(k) = V_{\mathfrak{a}}$. Hence $\Phi_{\mathfrak{f}, \mathfrak{a}}(x) = 1$ if $x \in V_{\mathfrak{a}} \cap V_{k, L_T}$ and 0 otherwise. Since $G(k) \cap G(k_\infty)G(\widehat{\mathcal{O}})_{\mathfrak{a}} = G_{\mathfrak{a}}$, we have

$$Z_{\mathfrak{a}}(\Phi, s) = \frac{1}{N(\mathfrak{a})^{2s}} \int_{G(k_\infty)/G_{\mathfrak{a}}} |\det g_\infty|_\infty^{2s} \sum_{x \in V_{\mathfrak{a}} \cap V_{L_T}} \Phi_\infty(g_\infty x) dg_\infty \cdot \int_{G(\widehat{\mathcal{O}})_{\mathfrak{a}}} dg_{\mathfrak{f}}.$$

Since $G(\mathbb{A}_{\mathfrak{f}})$ is unimodular, $\int_{G(\widehat{\mathcal{O}})_{\mathfrak{a}}} dg_{\mathfrak{f}} = 1$. Now the formula is obtained by the usual unfolding method. Note that $|\det g_\infty|_\infty^2 = |P(g_\infty x)/P(x)|_\infty$. \square

Combined with Lemma 4.2, we have the following.

Proposition 4.6.

$$Z(\Phi, s, \omega) = \mathcal{Z}_{L_T}(\Phi_\infty, s) \sum_{\mathfrak{a} \in \text{Cl}(k)} \omega(\mathfrak{a}) \vartheta_{L_T}(\mathfrak{a}, s).$$

We next discuss the contributions of irreducible and reducible forms.

Definition 4.7. We put

$$\begin{aligned} Z^{\text{ird}}(\Phi, s, \omega) &= \int_{G(\mathbb{A})/G(k)} |\det g|^{2s} \tilde{\omega}(\det g) \sum_{x \in V^{ss}(k)^{\text{ird}}} \Phi(gx) d_{\text{pr}} g, \\ \vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s) &= \sum_{R \in \mathcal{C}(\mathcal{O}, \mathfrak{a}, L_T)^{\text{ird}}} \frac{(\# \text{Aut}(R))^{-1}}{N(\Delta_R/\mathcal{O})^s}. \end{aligned}$$

We define $Z^{\text{rd}}(\Phi, s, \omega)$ and $\vartheta_{L_T}^{\text{rd}}(\mathfrak{a}, s)$ similarly.

By Proposition 3.5, the same argument of the proof of Lemma 4.5 shows the following.

Proposition 4.8.

$$\begin{aligned} Z^{\text{ird}}(\Phi, s, \omega) &= \mathcal{Z}_{L_T}(\Phi_\infty, s) \sum_{\mathfrak{a} \in \text{Cl}(k)} \omega(\mathfrak{a}) \vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s), \\ Z^{\text{rd}}(\Phi, s, \omega) &= \mathcal{Z}_{L_T}(\Phi_\infty, s) \sum_{\mathfrak{a} \in \text{Cl}(k)} \omega(\mathfrak{a}) \vartheta_{L_T}^{\text{rd}}(\mathfrak{a}, s). \end{aligned}$$

5. ANALYTIC CONTINUATIONS AND RESIDUE FORMULAE

In this section we give proofs of Theorem 1.1 (1) and Theorem 1.3 (1) with the values \mathfrak{A}_{L_T} , \mathfrak{B}_{L_T} in Theorem 1.2. We also give the residue of $\vartheta_{L_T}^{\text{rd}}(\mathfrak{a}, s)$ at $s = 1$.

We recall the residues of $Z^{\text{ird}}(\Phi, s, \omega)$ and $Z^{\text{rd}}(\Phi, s, \omega)$. For a while $\Phi \in \mathcal{S}(V(\mathbb{A}))$ and $\Phi_v \in \mathcal{S}(V(k_v))$ is arbitrary. We introduce the following distributions.

Definition 5.1. For $\Phi \in \mathcal{S}(V(\mathbb{A}))$, we define

$$\begin{aligned}\mathcal{A}(\Phi) &= \int_{\mathbb{A}^4} \Phi(x_0, x_1, x_2, x_3) d_{\text{pr}} x_0 d_{\text{pr}} x_1 d_{\text{pr}} x_2 d_{\text{pr}} x_3, \\ \mathcal{B}(\Phi) &= \int_{\mathbb{A}^\times \times \mathbb{A}^3} |t|_{\mathbb{A}}^{1/3} \Phi(t, x_1, x_2, x_3) d_{\text{pr}}^\times t d_{\text{pr}} x_1 d_{\text{pr}} x_2 d_{\text{pr}} x_3, \\ \mathcal{C}(\Phi) &= \int_{\mathbb{A}^\times \times \mathbb{A}^2} |t|_{\mathbb{A}}^2 \Phi(0, t, x_2, x_3) d_{\text{pr}}^\times t d_{\text{pr}} x_2 d_{\text{pr}} x_3.\end{aligned}$$

We have to give a notice for the definition of $\mathcal{B}(\Phi)$, because the integral defining $\mathcal{B}(\Phi)$ itself does not converge. (The integrals defining $\mathcal{A}(\Phi)$ and $\mathcal{C}(\Phi)$ converge.) From the Iwasawa-Tate theory, as a complex function of s the integral

$$\mathcal{B}(\Phi, s) = \int_{\mathbb{A}^\times \times \mathbb{A}^3} |t|_{\mathbb{A}}^s \Phi(t, x_1, x_2, x_3) d_{\text{pr}}^\times t d_{\text{pr}} x_1 d_{\text{pr}} x_2 d_{\text{pr}} x_3$$

has meromorphic continuation to the whole complex plane and is holomorphic except for $s = 0, 1$. The above definition means, more precisely speaking, we put $\mathcal{B}(\Phi) = \mathcal{B}(\Phi, 1/3)$.

The following is proved in [T06, Section 8].

Theorem 5.2. *Let Φ be \mathcal{K} -invariant. The integrals $Z^{\text{ird}}(\Phi, s, \omega)$ and $Z^{\text{rd}}(\Phi, s, \omega)$ have meromorphic continuations to the whole complex plane. Functions $(s-1)(s-5/6)Z^{\text{ird}}(\Phi, s, \omega)$ and $(s-1)Z^{\text{rd}}(\Phi, s, \omega)$ are holomorphic for $\text{Re}(s) > 1/2$, and the residues of $Z^{\text{ird}}(\Phi, s, \omega)$ and $Z^{\text{rd}}(\Phi, s, \omega)$ in this domain are given by the following;*

$$\begin{aligned}\text{Res}_{s=1} Z^{\text{ird}}(\Phi, s, \omega) &= \delta(\omega) 2^{-1} \pi^{-r_1} (2\pi)^{-r_2} \mathbf{c}_k \zeta_k(2) \mathcal{A}(\Phi), \\ \text{Res}_{s=5/6} Z^{\text{ird}}(\Phi, s, \omega) &= \delta(\omega^3) 6^{-1} \Delta_k^{-1/2} \mathbf{c}_k \mathcal{B}(\Phi), \\ \text{Res}_{s=1} Z^{\text{rd}}(\Phi, s, \omega) &= \delta(\omega) 2^{-1} \mathbf{c}_k \mathcal{C}(\Phi).\end{aligned}$$

We describe residues of $\vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s)$ and $\vartheta_{L_T}^{\text{rd}}(\mathfrak{a}, s)$ using this theorem and Proposition 4.8.

Definition 5.3. For $\Phi_v \in \mathcal{S}(V(k_v))$, we define

$$\begin{aligned}\mathcal{A}_v(\Phi_v) &= \int_{k_v^4} \Phi_v(x_0, x_1, x_2, x_3) dx_0 dx_1 dx_2 dx_3, \\ \mathcal{B}_v(\Phi_v) &= \int_{k_v^\times \times k_v^3} |t|_v^{1/3} \Phi_v(t, x_1, x_2, x_3) d^\times t dx_1 dx_2 dx_3, \\ \mathcal{C}_v(\Phi_v) &= \int_{k_v^\times \times k_v^2} |t|_v^2 \Phi_v(0, t, x_2, x_3) d^\times t dx_2 dx_3.\end{aligned}$$

Also we put

$$\mathcal{Z}_{L_v}(\Phi_v, s) = \int_{G(k_v)} |P(g_v x)|_v^s \Phi_v(g_v x) dg_v \quad (x \in V_{L_v}).$$

We now $\Phi = \prod_v \Phi_v$ is of the form in Assumption 4.3.

Definition 5.4. Let $\Phi_v \in \mathcal{S}(V(k_v))$ be as in Assumption 4.3.

(1) For $v \in \mathfrak{M}_\infty$, we define

$$\alpha_v(L_v) = \frac{\mathcal{A}_v(\Phi_v)}{\mathcal{Z}_{L_v}(\Phi_v, 1)}, \quad \beta_v(L_v) = \frac{\mathcal{B}_v(\Phi_v)}{\mathcal{Z}_{L_v}(\Phi_v, 5/6)}, \quad \gamma_v(L_v) = \frac{\mathcal{C}_v(\Phi_v)}{\mathcal{Z}_{L_v}(\Phi_v, 1)}$$

where Φ_v is chosen such that denominators do not vanish. It is known that these do not depend on the choice of such Φ_v .

(2) For $v \in \mathfrak{M}_f \cap T$, let $\Phi_{v,0}$ be the characteristic function of $V(\mathcal{O}_v)$. We put

$$\alpha_v(L_v) = \frac{\mathcal{A}_v(\Phi_v)}{\mathcal{A}_v(\Phi_{v,0})}, \quad \beta_v(L_v) = \frac{\mathcal{B}_v(\Phi_v)}{\mathcal{B}_v(\Phi_{v,0})}, \quad \gamma_v(L_v) = \frac{\mathcal{C}_v(\Phi_v)}{\mathcal{C}_v(\Phi_{v,0})}.$$

(3) Further we let the product as follows;

$$\alpha_T(L_T) = \prod_{v \in T} \alpha_v(L_v), \quad \beta_T(L_T) = \prod_{v \in T} \beta_v(L_v), \quad \gamma_T(L_T) = \prod_{v \in T} \gamma_v(L_v).$$

Theorem 5.5. (1) *The Dirichlet series $\vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s)$ and $\vartheta_{L_T}^{\text{rd}}(\mathbf{a}, s)$ have meromorphic continuations to the whole complex plane. Functions $(s-1)(s-5/6)\vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s)$ and $(s-1)\vartheta_{L_T}^{\text{rd}}(\mathbf{a}, s)$ are holomorphic for $\text{Re}(s) > 1/2$.*

(2) *The residues of $\vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s)$ and $\vartheta_{L_T}^{\text{rd}}(\mathbf{a}, s)$ in the region $\text{Re}(s) > 1/2$ are given by;*

$$\begin{aligned} \text{Res}_{s=1} \vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s) &= h_k^{-1} 2^{-1} \pi^{-r_1} (2\pi)^{-r_2} \mathbf{c}_k \zeta_k(2) \alpha_T(L_T), \\ \text{Res}_{s=5/6} \vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s) &= \tau(\mathbf{a}) h_k^{(3)} h_k^{-1} 6^{-1} \Delta_k^{-1/2} \mathbf{c}_k \zeta_k(1/3) \beta_T(L_T), \\ \text{Res}_{s=1} \vartheta_{L_T}^{\text{rd}}(\mathbf{a}, s) &= h_k^{-1} 2^{-1} \mathbf{c}_k \zeta_k(2) \gamma_T(L_T). \end{aligned}$$

Proof. By the theory of prehomogeneous vector spaces, the local zeta function $\mathcal{Z}_{L_T}(\Phi_\infty, s)$ has meromorphic continuation to the whole complex plane. Moreover, for any $s \in \mathbb{C}$ such that $\text{Re}(s) > 1/6$, we can choose Φ_∞ such that $\mathcal{Z}_{L_T}(\Phi_\infty, s) \neq 0$. Hence (1) follows from Proposition 4.8, Theorem 5.2 and the orthogonality of characters.

We consider (2). We prove the second formula. The first and third formulae are proved similarly. Let $\vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s, \omega) = \sum_{\mathbf{a} \in \text{Cl}(k)} \omega(\mathbf{a}) \vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s)$, which is the Dirichlet series appeared in the first formula of Proposition 4.8. For $\Psi_f \in \mathcal{S}(V(\mathbb{A}_f))$ we put the “finite part” $\mathcal{B}_f(\Psi_f)$ of \mathcal{B} by evaluating at $s = 1/3$ of the analytic function $\int_{\mathbb{A}_f^\times \times \mathbb{A}_f^3} |t|_{\mathbb{A}_f}^s \Psi_f(t, x_1, x_2, x_3) d_f^\times t d_f x_1 d_f x_2 d_f x_3$. Then $\mathcal{B}(\Phi) = \mathcal{B}_f(\Phi_f) \prod_{v \in \mathfrak{M}_\infty} \mathcal{B}_v(\Phi_v)$. Let $\Phi_{f,0}$ be the characteristic function of $V(\widehat{\mathcal{O}})$. Then we have $\mathcal{B}(\Phi_{f,0}) = \zeta_k(1/3)$ and $\mathcal{B}_f(\Phi_f)/\mathcal{B}_f(\Phi_{f,0}) = \prod_{v \in T \cap \mathfrak{M}_f} \mathcal{B}_v(\Phi_v)/\mathcal{B}_v(\Phi_{v,0})$ by the analytic continuations. Hence by Proposition 4.8 and Theorem 5.2 we have

$$\begin{aligned} \text{Res}_{s=5/6} \vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s, \omega) &= \delta(\omega^3) \frac{\mathbf{c}_k \zeta_k(1/3)}{6 \Delta_k^{1/2}} \frac{\prod_{v \in \mathfrak{M}_\infty} \mathcal{B}_v(\Phi_v)}{\mathcal{Z}_{L_T}(\Phi_\infty, 5/6)} \prod_{v \in T \cap \mathfrak{M}_f} \frac{\mathcal{B}_v(\Phi_v)}{\mathcal{B}_v(\Phi_{v,0})} \\ &= \delta(\omega^3) 6^{-1} \Delta_k^{-1/2} \mathbf{c}_k \zeta_k(1/3) \beta_T(L_T). \end{aligned}$$

By the orthogonality of the characters, $\vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s) = \sum_{\omega \in \text{Cl}(k)^*} \omega(\mathbf{a})^{-1} \vartheta_{L_T}^{\text{ird}}(\mathbf{a}, s, \omega)$. From the identity $\sum_{\omega \in \text{Cl}(k)^*} \omega(\mathbf{a})^{-1} \delta(\omega^3) = \tau(\mathbf{a}) h_k^{(3)} / h_k$, we have the second formula of (2). \square

From now on we consider $\alpha_v(L_v)$, $\beta_v(L_v)$, and $\gamma_v(L_v)$. For $v \in \mathfrak{M}_\infty$, this is done by Shintani [S72] and Datskovsky-Wright [DW86] and we already used their result in [T06]. For the convenience of the reader, we give these values in Table 1:

Proposition 5.6. *For $v \in \mathfrak{M}_\infty$, $\alpha_v(L_v)$, $\beta_v(L_v)$ and $\gamma_v(L_v)$ are given by Table 1.*

We compute $\alpha_v(L_v)$, $\beta_v(L_v)$, and $\gamma_v(L_v)$ for $v \in \mathfrak{M}_f$ using the local theory developed by Datskovsky-Wright [DW86], especially Theorems 3.1, 5.1, 5.2 and Propositions 5.1, 5.3. We fix $v \in \mathfrak{M}_f$ for the rest of this section. To simplify the notation, we drop the subscript v and write L , q , α , β , γ , \mathcal{A} , \mathcal{B} and \mathcal{C} instead of writing L_v , q_v , α_v , β_v , γ_v , \mathcal{A}_v , \mathcal{B}_v and \mathcal{C}_v . To stress

| v | L_v | $\alpha_v(L_v)$ | $\beta_v(L_v)$ | $\gamma_v(L_v)$ |
|--------------|--|------------------|--|-----------------|
| \mathbb{R} | $\mathbb{R} \times \mathbb{R} \times \mathbb{R}$ | $\frac{\pi}{6}$ | $\frac{\sqrt{3}\Gamma(1/3)^3}{4\pi}$ | $\frac{1}{2}$ |
| \mathbb{R} | $\mathbb{R} \times \mathbb{C}$ | $\frac{\pi}{2}$ | $\frac{3\Gamma(1/3)^3}{4\pi}$ | $\frac{1}{2}$ |
| \mathbb{C} | $\mathbb{C} \times \mathbb{C} \times \mathbb{C}$ | $\frac{2\pi}{6}$ | $\frac{\sqrt{3}\Gamma(1/3)^6}{8\pi^2}$ | $\frac{1}{2}$ |

TABLE 1.

the dependence Φ_v in Assumption 4.3 on L , we write $\Phi_v = \Phi_L$. As in Section 1, we denote by θ_L the order of the automorphisms of L as a k_v -algebra. Let $\zeta_L(s)$ be the local zeta function associated with the k_v -algebra L . Namely, we define

$$\zeta_L(s) = \begin{cases} (1 - q^{-s})^{-3} & L = k_v \times k_v \times k_v, \\ (1 - q^{-s})^{-1}(1 - q^{-2s})^{-1} & L = k_v \times (\text{quad. unramified ext. of } k_v), \\ (1 - q^{-3s})^{-1} & L = (\text{cubic unramified ext. of } k_v), \\ (1 - q^{-s})^{-2} & L = k_v \times (\text{quad. ramified ext. of } k_v), \\ (1 - q^{-s})^{-1} & L = (\text{cubic ramified ext. of } k_v). \end{cases}$$

Let Δ_{L/k_v} be the discriminant of L/k_v and Δ_L its norm. An element x_L in the $G(k_v)$ -orbit V_L is called a *standard orbital representative* if $x_L \in V(\mathcal{O}_v)$ and $P(x_L) \in \mathcal{O}_v$ generates the ideal Δ_{L/k_v} . It is easy to see that such an element exists for arbitrary L . We fix such x_L . We define

$$\Omega_L(\Phi_L, s) = \int_{G(k_v)} |\det g_v|_v^{2s} \Phi_L(g_v x_L) dg_v,$$

which equals to $|P(x_L)|_v^{-s} \mathcal{Z}_{L_v}(\Phi_v, s)$ and hence do not depend on the choice of x_L . This function plays an important role in the computation.

Before starting the computation, we will compare and adjust the notation in [DW86] to ours. In [DW86] they denote by A the set of $G(k_v)$ -orbits of $V^{\text{ss}}(k_v)$ and by α any of its element. Hence there is the canonical correspondence between their A and our $\mathcal{C}^{\text{sep}}(k_v)$. If $L \in \mathcal{C}^{\text{sep}}(k_v)$ corresponds to $\alpha \in A$, then $\Omega_L(\Phi_L, s)$ equals to what they denoted by $Z_\alpha(\omega_{2s}, \Phi_L)$. (They defined the distribution $Z_\alpha(\omega, \Phi)$ in p.39.) Also the value c_α they introduced in p.38 equals to $\theta_L^{-1} \Delta_L^{-1}$, since their $o(\alpha)$ equals to θ_L and their x_α is also a standard orbital representative for the orbit V_L .

The following beautiful formula is a variation of [DW86, Theorem 3.1].

Lemma 5.7.

$$\Omega_L(\Phi_L, s) = (1 - q^{-4s})^{-1} (1 - q^{-6s+1})^{-1} \zeta_L(2s) \zeta_L(4s)^{-1}.$$

Proof. Let $\Phi_0 \in \mathcal{A}(V(k_v))$ be the characteristic function of $V(\mathcal{O}_v)$. Then since Φ_L is the characteristic function of $V(\mathcal{O}_v) \cap G(k_v)x_L$, $\Phi_L(g_v x_L) = \Phi_0(g_v x_L)$ for all $g_v \in G(k_v)$. Hence

$$\Omega_L(\Phi_L, s) = \int_{G(k_v)} |\det g_v|_v^{2s} \Phi_0(g_v x_L) dg_v.$$

The right hand side is what Datskovsky and Wright gave the explicit formula in [DW86, Theorem 3.1]. Since their notation $\omega(\pi)$ is q^{-2s} in our setting, we have the formula. \square

Proposition 5.8.

$$\alpha(L) = \frac{(1 - q^{-1})(1 - q^{-2})}{(1 - q^{-4})(1 - q^{-5})} \cdot \theta_L^{-1} \Delta_L^{-1} \cdot \frac{\zeta_L(2)}{\zeta_L(4)}.$$

Proof. Obviously $\mathcal{A}(\Phi_0) = 1$ and hence $\alpha(L) = \mathcal{A}(\Phi_L)$. This is expressed by $q^{2e}\widehat{\Phi}_L(0)$ in the notation of [DW86]. Note that their additive measure on k_v is $q^{-e/2}$ times to ours. Since $\Phi_L(x) = 0$ if $x \notin V_L$, by [DW86, Propostion 5.1] we have

$$\mathcal{A}(\Phi_L) = (1 - q^{-1})(1 - q^{-2})\theta_L^{-1}\Delta_L^{-1}\Omega_L(\Phi_L, 1).$$

Hence from Lemma 5.7 we have the formula. \square

Proposition 5.9.

$$\beta(L) = \frac{(1 - q^{-1/3})(1 - q^{-1})}{(1 - q^{-10/3})(1 - q^{-4})} \cdot \theta_L^{-1}\Delta_L^{-1} \cdot \frac{\zeta_L(1/3)\zeta_L(5/3)}{\zeta_L(2/3)\zeta_L(10/3)}.$$

Proof. Since $\mathcal{B}(\Phi_0) = (1 - q^{-1/3})^{-1}$ we have $\beta(L) = (1 - q^{-1/3})\mathcal{B}(\Phi_L)$. If we use the notation of [DW86], $\mathcal{B}(\Phi_L)$ equals to $q^{3e/2}\Sigma_4(1, \Phi_L)$ where 1 denotes the trivial character on k_v . Hence if L corresponds to $\alpha \in A$ in their notation, by [DW86, Theorem 5.2] we have

$$\mathcal{B}(\Phi_L) = (1 - q^{-1})\theta_L^{-1}\Delta_L^{-1}a_\alpha(1)\Omega_L(\Phi_L, 5/6).$$

The value $a_\alpha(1)$ is evaluated in [DW86, Proposition 5.3] and is found to be $\zeta_L(1/3)\zeta_L(2/3)^{-1}$. Now Lemma 5.7 gives the desired formula. \square

Theorem 1.1 (1) and Theorem 1.3 (1) with the values $\mathfrak{A}_{L_T}, \mathfrak{B}_{L_T}$ in Theorem 1.2 follows from Theorem 5.5 and Propositions 5.6, 5.8, 5.9. We also evaluate $\gamma(L)$. This gives the explicit formula of the residue of $\vartheta_{L_T}^{\text{rd}}(\mathfrak{a}, s)$ at $s = 1$.

Proposition 5.10.

$$\gamma(L) = \begin{cases} \frac{(1 - q^{-1})(1 - q^{-2})}{(1 - q^{-4})(1 - q^{-5})} \cdot 2^{-1}\Delta_L^{-1} \cdot \frac{\zeta_L(2)}{\zeta_L(4)} & L \text{ is not a field,} \\ 0 & L \text{ is a field.} \end{cases}$$

Proof. The value $\mathcal{C}(\Phi_L)$ equals to $q^e\Sigma_2(\Phi_L)$ if we use the notation of [DW86]. Hence by [DW86, Theorem 5.2], this equals to $(1 - q^{-1})2^{-1}\Delta_L^{-1}\Omega_L(\Phi_L, 1)$ if L is not a field and 0 otherwise. Since $\mathcal{C}(\Phi_0) = (1 - q^{-2})^{-1}$, we have the formula. \square

Remark 5.11. Let $\mathcal{E}_i^{\text{rm}}(k_v)$ the set of isomorphism classes of totally ramified extensions of k_v of degree i . For $F \in \mathcal{E}_i^{\text{rm}}(k_v)$, let Δ_F be the norm of the relative discriminant of F/k_v and θ_F the order of automorphisms of F . In [S78] Serre established a beautiful formula

$$\sum_{F \in \mathcal{E}_i^{\text{rm}}(k_v)} \theta_F^{-1}\Delta_F^{-1} = q^{-i+1}.$$

On the other side, by gathering all the splitting type at a fixed $v \in \mathfrak{M}_f$, we have

$$\sum_{L \in \mathcal{C}^{\text{sep}}(k_v)} \alpha(L) = \sum_{L \in \mathcal{C}^{\text{sep}}(k_v)} \beta(L) = \sum_{L \in \mathcal{C}^{\text{sep}}(k_v)} \gamma(L) = 1.$$

By computation we see that this is equivalent to the Serre's formula for $i = 2$ and 3.

6. DENSITY THEOREMS

In this section we give proofs of Theorem 1.1 (2) and Theorem 1.3 (2). Throughout this section we fix L_T . Recall that we define Dirichlet series $\vartheta_{L_T}(\mathfrak{a}, s)$, $\vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s)$, $\vartheta_{L_T}^{\text{rd}}(\mathfrak{a}, s)$ in Section 4. These satisfy $\vartheta_{L_T}(\mathfrak{a}, s) = \vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s) + \vartheta_{L_T}^{\text{rd}}(\mathfrak{a}, s)$. We put $\vartheta_{L_T}(s) = \sum_{\mathfrak{a} \in \text{Cl}(k)} \vartheta_{L_T}(\mathfrak{a}, s)$. Similarly we define $\vartheta_{L_T}^{\text{ird}}(s)$, $\vartheta_{L_T}^{\text{rd}}(s)$ as the sum of the Dirichlet series over $\mathfrak{a} \in \text{Cl}(k)$. We write

$$\vartheta_{L_T}(s) = \sum_{m \geq 1} a_m/m^s, \quad \vartheta_{L_T}^{\text{ird}}(s) = \sum_{m \geq 1} a_m^{\text{ird}}/m^s, \quad \vartheta_{L_T}^{\text{rd}}(s) = \sum_{m \geq 1} a_m^{\text{rd}}/m^s,$$

so that $a_n = a_n^{\text{ird}} + a_n^{\text{rd}}$. Until the proof of Lemma 6.5 we prove the following.

Proposition 6.1. *For any $\varepsilon > 0$,*

$$\sum_{m < X} a_m^{\text{ird}} = \mathfrak{A}_{L_T} X + (5/6)^{-1} \mathfrak{B}_{L_T} X^{5/6} + O(X^{\frac{5n-1}{5n+1} + \varepsilon}) \quad (X \rightarrow \infty).$$

To prove the proposition we first give an estimate of the function $\sum_{m < X} a_m$. We put $\mathfrak{C}_{L_T} = 2^{-1} \mathfrak{c}_k \zeta_k(2) \prod_{v \in T} \gamma_v(L_v)$, which is the residue of $\vartheta_{L_T}^{\text{rd}}(s)$ at $s = 1$.

Lemma 6.2. *For any $\varepsilon > 0$,*

$$\sum_{m < X} a_m = (\mathfrak{A}_{L_T} + \mathfrak{C}_{L_T}) X + (5/6)^{-1} \mathfrak{B}_{L_T} X^{5/6} + O(X^{\frac{4n-1}{4n+1} + \varepsilon}) \quad (X \rightarrow \infty).$$

Proof. In [W85] Wright proved that the global zeta function $Z(\Phi, s, \omega)$ can be continued holomorphically to the whole complex plane except for possible simple poles at $s = 0, 1/6, 5/6, 1$ and satisfies the functional equation

$$Z(\Phi, s, \omega) = Z(\hat{\Phi}, 1 - s, \omega^{-1})$$

where $\hat{\Phi}$ is a suitable Fourier transform of Φ . Let $\omega \in \text{Cl}(k)^*$ be the trivial character and $\Phi \in \mathcal{S}(V(\mathbb{A}))$ as in Assumption 4.3. Then by Proposition 4.6 $Z(\Phi, s, \omega) = \mathcal{Z}_{L_T}(\Phi_\infty, s) \vartheta_{L_T}(s)$. Since $\hat{\Phi}$ is also a $G(\hat{\mathcal{O}})$ -invariant function, we have a similar decomposition for $Z(\hat{\Phi}, s, \omega^{-1})$. These combined with archimedean local theory show that $\vartheta_{L_T}(s)$ is holomorphic except for $s = 1, 5/6$ and satisfies a functional equation of the form

$$\vartheta_{L_T}(1 - s) = \Gamma(s)^{2n} \Gamma(s - \frac{1}{6})^n \Gamma(s + \frac{1}{6})^n \sum_{\lambda \in \Lambda} p_\lambda(e^{\pi\sqrt{-1}s/2}, e^{-\pi\sqrt{-1}s/2}) \xi_\lambda(s)$$

where Λ is a finite index set and for each $\lambda \in \Lambda$, $p_\lambda(x, y)$ is a polynomial in x, y of degree less than $4n$ and $\xi_\lambda(s)$ is a Dirichlet series with absolute convergence domain $\text{Re}(s) > 1$. By Theorem 5.5 the residues of $\vartheta_{L_T}(s)$ at $s = 1$ and $5/6$ are $\mathfrak{A}_{L_T} + \mathfrak{C}_{L_T}$ and \mathfrak{B}_{L_T} , respectively. Hence the proposition follows from the modified Landau theorem [SS74, Theorem 3]. \square

This lemma reduces the proof of Proposition 6.1 to an estimate of $\sum_{m < X} a_m^{\text{rd}}$. To give an estimate of this function we introduce another prehomogeneous vector space. Let $B = B_2 \subset G$ and W the subspace of V having a linear factor v ;

$$B = \left\{ \begin{pmatrix} * & 0 \\ * & * \end{pmatrix} \right\}, \quad W = \{y = y(u, v) = v(y_1 u^2 + y_2 uv + y_3 v^2) \mid y_1, y_2, y_3 \in \text{Aff}\}.$$

Then W is an invariant subspace of B and (B, W) is also a prehomogeneous vector space. As in [T06], we use the functional equation of the zeta function for this space to study $\sum_{m < X} a_m^{\text{rd}}$. (This idea is due to Shintani [S75].) For $\mathfrak{a}, \mathfrak{c} \in \text{Cl}(k)$, we put

$$B(k) \supset B_{\mathfrak{a}, \mathfrak{c}} = \left\{ \begin{pmatrix} t & 0 \\ u & p \end{pmatrix} \mid t, p \in \mathcal{O}^\times, u \in \mathfrak{a}^{-1} \mathfrak{c}^{-2} \right\},$$

$$W(k) \supset W_{\mathfrak{a}, \mathfrak{c}} = \{y \mid y_1 \in \mathfrak{c}, y_2 \in \mathfrak{a}^{-1} \mathfrak{c}^{-1}, y_3 \in \mathfrak{a}^{-2} \mathfrak{c}^{-3}\}.$$

Then $W_{\mathfrak{a}, \mathfrak{c}}$ is $B_{\mathfrak{a}, \mathfrak{c}}$ -invariant. Let $W_{k, L_T} = W(k) \cap V_{k, L_T}$. We define

$$\eta_{L_T}(s) = \sum_{\mathfrak{a}, \mathfrak{c} \in \text{Cl}(k)} \sum_{y \in B_{\mathfrak{a}, \mathfrak{c}} \setminus (W_{\mathfrak{a}, \mathfrak{c}} \cap W_{k, L_T})} \frac{(\# \text{Stab}(B_{\mathfrak{a}, \mathfrak{c}}; y))^{-1}}{N(\mathfrak{a})^{2s} |P(y)|_\infty^s}.$$

We write $\eta_{L_T}(s) = \sum_{m \geq 1} b_m / m^s$. Note that for $y \in W_{\mathfrak{a}, \mathfrak{c}}$, $P(y) \in \mathfrak{a}^{-2}$ and hence $N(\mathfrak{a})^2 |P(y)|_\infty$ is an integer. Then we have the following.

Lemma 6.3. *For any $\varepsilon > 0$,*

$$\sum_{m < X} b_m = \mathfrak{C}_{L_T} X + O(X^{\frac{5n-1}{5n+1} + \varepsilon}) \quad (X \rightarrow \infty).$$

Proof. We give an integral expression of $\eta_{L_T}(s)$. For $\Psi \in \mathcal{S}(W(\mathbb{A}))$ and $s \in \mathbb{C}$, define

$$Y(\Psi, s) = \int_{B(\mathbb{A})/B(k)} |\det b|_{\mathbb{A}}^{2s} \sum_{y \in W(k) \cap V^{\text{ss}}(k)} \Psi(by) db.$$

Also for $\Psi_{\infty} \in \mathcal{S}(W(k_{\infty}))$, we put

$$\mathcal{Y}_{L_T}(\Psi_{\infty}, s) = \int_{B(k_{\infty})} |P(b_{\infty}y)|_{\infty}^s \Psi_{\infty}(b_{\infty}y) db_{\infty} \quad (y \in W_{k, L_T}).$$

Let $\Phi \in \mathcal{S}(V(\mathbb{A}))$ be as in Assumption 4.3. We regard $\Phi \in \mathcal{S}(W(\mathbb{A}))$ via the pullback of the inclusion $W(\mathbb{A}) \rightarrow V(\mathbb{A})$. Then since

$$\begin{aligned} B(\mathbb{A}) &= \coprod_{\mathfrak{a}, \mathfrak{c} \in \text{Cl}(k)} B(k_{\infty})B(\widehat{\mathcal{O}}) \cdot \text{diag}(\mathfrak{c}, \mathfrak{a}^{-1}\mathfrak{c}^{-1}) \cdot B(k), \\ B_{\mathfrak{a}, \mathfrak{c}} &= B(k) \cap \text{diag}(\mathfrak{c}, \mathfrak{a}^{-1}\mathfrak{c}^{-1})^{-1} \cdot B(k_{\infty})B(\widehat{\mathcal{O}}) \cdot \text{diag}(\mathfrak{c}, \mathfrak{a}^{-1}\mathfrak{c}^{-1}), \\ W_{\mathfrak{a}, \mathfrak{c}} &= W(k) \cap (\text{diag}(\mathfrak{c}, \mathfrak{a}^{-1}\mathfrak{c}^{-1})^{-1}W(\widehat{\mathcal{O}}) \times W(k_{\infty})), \end{aligned}$$

by the similar unfolding method as in Proposition 4.6 we have

$$Y(\Phi, s) = \mathcal{Y}_{L_T}(\Phi_{\infty}, s)\eta_{L_T}(s).$$

We recall from [T06] some results on zeta functions for (B, W) . The residue of $Y(\Phi, s)$ at $s = 1$ is exactly the same as that of $Z^{\text{rd}}(\Phi, s)$. Also since Φ_{∞} is $\prod_{v \in \mathfrak{M}_{\infty}} \mathcal{K}_v$ -invariant, the local zeta functions $\mathcal{Z}_{L_T}(\Phi_{\infty}, s)$ and $\mathcal{Y}_{L_T}(\Phi_{\infty}, s)$ also coincide. Hence the residue of $\eta_{L_T}(s)$ at $s = 1$ is \mathfrak{C}_{L_T} . Moreover, the Dirichlet series $H(s) = \eta_{L_T}(s)\zeta_k(4s)$ is entire after multiplied by $(s - 1/2)^3(s - 1)$, and it satisfies a functional equation of the form

$$H(1 - s) = \Gamma(2s - 1)^{2n} \Gamma(s)^n \Gamma(3s - 3/2)^n \sum_{\lambda \in \Lambda} q_{\lambda}(e^{\pi\sqrt{-1}s/2}, e^{-\pi\sqrt{-1}s/2}) \theta_{\lambda}(s)$$

where Λ is a finite index set and for each $\lambda \in \Lambda$, $q_{\lambda}(x, y)$ is a polynomial in x, y of degree less than $8n$ and $\theta_{\lambda}(s)$ is a Dirichlet series with absolute convergence domain $\text{Re}(s) > 1$. Hence by the same argument as in [T06, Proposition 7.18] we have the lemma. \square

We introduce the following notation.

Definition 6.4. Let $k(s) = \sum_{m \geq 1} k_m/m^s$ and $l(s) = \sum_{m \geq 1} l_m/m^s$ be Dirichlet series having absolute convergence domains. We say $k(s)$ is bounded by $l(s)$ if $|k_m| \leq l_m$ for all $m \geq 1$, and write $k(s) \preceq l(s)$ in this situation.

Using Lemma 6.3 we have the following estimate, which proves Proposition 6.1. Notice that for a positive sequence $\{a_m\}$ and a positive constant ρ , the series $\sum_{m \geq 1} a_m/m^s$ converges for $\text{Re}(s) > \rho$ if and only if $\sum_{m < X} a_m = O(X^{\rho+\epsilon})$ for any $\epsilon > 0$.

Lemma 6.5. For any $\varepsilon > 0$,

$$\sum_{m < X} a_m^{\text{rd}} = \mathfrak{C}_{L_T} X + O(X^{\frac{5n-1}{5n+1}+\varepsilon}) \quad (X \rightarrow \infty).$$

Proof. Let $V_k^{(1)} = \{x \in V^{\text{ss}}(k) \mid k(x) = k \times k \times k\}$, $V_k^{(2)} = V^{\text{ss}}(k) \setminus (V^{\text{ss}}(k)^{\text{ird}} \cup V_k^{(1)})$ and put $V_{\mathfrak{a}}^{(i)} = V_{\mathfrak{a}} \cap V_k^{(i)}$, $W_{\mathfrak{a}, \mathfrak{c}}^{(i)} = W_{\mathfrak{a}, \mathfrak{c}} \cap V_k^{(i)}$ for $i = 1, 2$. In [T06, Proposition 3.12], we construct a bijective map

$$\psi_{\mathfrak{a}}: \coprod_{\mathfrak{c} \in \text{Cl}(k)} (B_{\mathfrak{a}, \mathfrak{c}} \setminus W_{\mathfrak{a}, \mathfrak{c}}^{(2)}) \longrightarrow G_{\mathfrak{a}} \setminus V_{\mathfrak{a}}^{(2)}$$

which preserves the value of P up to $(\mathcal{O}^\times)^2$ -multiple and $\text{Stab}(B_{\mathfrak{a},\mathfrak{c}}; y) \cong \text{Stab}(G_{\mathfrak{a}}; \psi_{\mathfrak{a}}(y))$ for any $y \in W_{\mathfrak{a},\mathfrak{c}}^{(2)}$. From the construction it is easy to see that this induces a bijective map

$$\psi_{\mathfrak{a},L_T}: \coprod_{\mathfrak{c} \in \text{Cl}(k)} \left(B_{\mathfrak{a},\mathfrak{c}} \setminus (W_{\mathfrak{a},\mathfrak{c}}^{(2)} \cap W_{k,L_T}) \right) \longrightarrow G_{\mathfrak{a}} \setminus (V_{\mathfrak{a}}^{(2)} \cap V_{k,L_T}).$$

On the other hand by the local theory, the Dirichlet series

$$\sum_{x \in G_{\mathfrak{a}} \setminus (V_{\mathfrak{a}}^{(1)} \cap V_{k,L_T})} N(\mathfrak{a})^{-2s} |P(x)|_{\infty}^{-s}, \quad \sum_{y \in B_{\mathfrak{a},\mathfrak{c}} \setminus (W_{\mathfrak{a},\mathfrak{c}}^{(1)} \cap W_{k,L_T})} N(\mathfrak{a})^{-2s} |P(y)|_{\infty}^{-s}$$

are bounded by the Dirichlet series $\zeta_k(2s)^3 \zeta_k(6s-1)/\zeta_k(4s)^2$. Especially these Dirichlet series converges for $\text{Re}(s) > 1/3$. Hence $\sum_{m < X} a_m^{\text{rd}} = \sum_{m < X} b_m + O(X^{1/3+\epsilon})$ where $\epsilon > 0$ is arbitrary and this finishes the proof. \square

We now give a proof of Theorem 1.1 (2).

Theorem 6.6. *Let $h_{L_T}(X)$ be as in Section 1. For any $\varepsilon > 0$,*

$$h_{L_T}(X) = \mathfrak{A}_{L_T} X + (5/6)^{-1} \mathfrak{B}_{L_T} X^{5/6} + O(X^{\frac{5n-1}{5n+1}+\varepsilon}) \quad (X \rightarrow \infty).$$

Proof. Let

$$\tilde{\vartheta}_{L_T}^{\text{ird}}(s) = \sum_{n \geq 1} \frac{\tilde{a}_n^{\text{ird}}}{n^s} = \sum_{R \in \mathcal{C}(\mathcal{O}, L_T)^{\text{ird}}} \frac{1}{N(\Delta_{R/\mathcal{O}})^s}.$$

Then $h_{L_T}(X) = \sum_{m < X} \tilde{a}_m^{\text{ird}}$. We compare $\vartheta_{L_T}^{\text{ird}}(s)$ and $\tilde{\vartheta}_{L_T}^{\text{ird}}(s)$. Let \mathfrak{G} be the set of isomorphism classes of cyclic cubic extensions of k . Then $\text{Aut}(R)$ is non-trivial only if $R \otimes k \in \mathfrak{G}$. Hence as in the proof of [T06, Theorem 7.20],

$$\begin{aligned} \tilde{\vartheta}_{L_T}^{\text{ird}}(s) - \vartheta_{L_T}^{\text{ird}}(s) &\leq \sum_{F \in \mathfrak{G}} \sum_{R \in \mathcal{C}(\mathcal{O}, R \otimes k = F)} \frac{1}{N(\Delta_{R/\mathcal{O}})^s} \\ &= \sum_{F \in \mathfrak{G}} N(\Delta_{F/k})^{-s} \zeta_k(4s) \zeta_k(6s-1) \zeta_F(2s) \zeta_F(4s)^{-1} \\ &\leq \zeta_k(2s)^3 \zeta_k(6s-1) \zeta_k(4s)^{-2} \sum_{F \in \mathfrak{G}} N(\Delta_{F/k})^{-s} \end{aligned}$$

and the Dirichlet series in the last term has absolute convergence domain $\text{Re}(s) > 1/2$. This shows that $h_{L_T}(X) = \sum_{m < X} \tilde{a}_m^{\text{ird}} = \sum_{m < X} a_m^{\text{ird}} + O(X^{1/2+\epsilon})$ where $\epsilon > 0$ is arbitrary. Hence from Proposition 6.1 we have the desired result. \square

We conclude this paper with a proof of Theorem 1.3 (2), which is much simpler than that of Theorem 6.6.

Theorem 6.7. *Assume L_T is chosen so that at least one of L_v is a field. Then $\vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s)$ is holomorphic in the whole complex plane except for $s = 1$ and $5/6$. Also for any $\varepsilon > 0$,*

$$h_{L_T}(\mathfrak{a}, X) = \frac{\mathfrak{A}_{L_T}}{h_k} X + \tau(\mathfrak{a}) \frac{h_k^{(3)}}{h_k} \cdot \frac{\mathfrak{B}_{L_T}}{5/6} X^{5/6} + O(X^{\frac{4n-1}{4n+1}+\varepsilon}) \quad (X \rightarrow \infty).$$

Proof. The essential fact that we can prove this formula is that under the condition on L_T we have $\mathcal{C}(\mathcal{O}, L_T)^{\text{ird}} = \mathcal{C}(\mathcal{O}, L_T)$. By the global and local theory for the space of binary cubic forms and the orthogonality of characters, $\vartheta_{L_T}^{\text{ird}}(\mathfrak{a}, s) = \vartheta_{L_T}(\mathfrak{a}, s)$ is holomorphic except for $s = 1, 5/6$ and satisfies a functional equation of the form

$$\vartheta_{L_T}(\mathfrak{a}, 1-s) = \Gamma(s)^{2n} \Gamma(s - \frac{1}{6})^n \Gamma(s + \frac{1}{6})^n \sum_{\lambda' \in \Lambda'} p_{\lambda'}(e^{\pi\sqrt{-1}s/2}, e^{-\pi\sqrt{-1}s/2}) \xi_{\lambda'}(s)$$

where Λ' is a finite index set and for each $\lambda' \in \Lambda'$, $p_{\lambda'}(x, y)$ is a polynomial in x, y of degree less than $4n$ and $\xi_{\lambda'}(s)$ is a Dirichlet series with absolute convergence domain $\operatorname{Re}(s) > 1$. Hence the theorem follows from the modified Landau theorem [SS74, Theorem 3], the residue formulae of $\vartheta_{L_T}(\mathfrak{a}, s)$ in Theorem 1.3 (1) and the same treatment for orders of cyclic Galois extensions as in the proof of Theorem 6.6. \square

REFERENCES

- [DW86] B. Datskovsky and D.J. Wright. The adelic zeta function associated with the space of binary cubic forms II: Local theory. *J. Reine Angew. Math.*, 367:27–75, 1986.
- [DW88] B. Datskovsky and D.J. Wright. Density of discriminants of cubic extensions. *J. Reine Angew. Math.*, 386:116–138, 1988.
- [DF64] B.N. Delone and Faddeev D.K. *The theory of irrationalities of the third degree*, volume 10 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, 1964.
- [L15] E. Landau. Über die Anzahl der Gitterpunkten in gewissen Bereichen II. *Gött. Nach.*, pages 209–243, 1915.
- [M71] J. Milnor. *Introduction to algebraic K-theory*. Princeton University Press, 1971.
- [SS74] M. Sato and T. Shintani. On zeta functions associated with prehomogeneous vector spaces. *Ann. of Math.*, 100:131–170, 1974.
- [S78] J.P. Serre. Une “formule de masse” pour les extensions totalement ramifiées de degré donné d’un corps local. *C. R. Acad. Sci. Paris A*, 286:1031–1036, 1978.
- [S72] T. Shintani. On Dirichlet series whose coefficients are class-numbers of integral binary cubic forms. *J. Math. Soc. Japan*, 24:132–188, 1972.
- [S75] T. Shintani. On zeta-functions associated with vector spaces of quadratic forms. *J. Fac. Sci. Univ. Tokyo, Sect IA*, 22:25–66, 1975.
- [T06] T. Taniguchi. Distributions of discriminants of cubic algebras. Preprint 2006, math.NT/0606109.
- [W85] D.J. Wright. The adelic zeta function associated to the space of binary cubic forms part I: Global theory. *Math. Ann.*, 270:503–534, 1985.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO
E-mail address: `tani@ms.u-tokyo.ac.jp`